arXiv:1703.06775v1 [math.FA] 20 Mar 2017

DENSITY OF TRANSLATES IN WEIGHTED L^p SPACES ON LOCALLY COMPACT GROUPS

EVGENY ABAKUMOV AND YULIA KUZNETSOVA

Abstract. Let G be a locally compact group, and let $1 \leqslant p < \infty$. Consider the weighted L^p-space $L^p(G,\omega) = \{f : \int |f\omega|^p < \infty\}$, where $\omega : G \to \mathbb{R}$ is a positive measurable function. Under appropriate conditions on ω, G acts on $L^p(G,\omega)$ by translations. When is this action hypercyclic, that is, there is a function in this space such that the set of all its translations is dense in $L^p(G,\omega)$? H. Salas (1995) gave a criterion of hypercyclicity in the case $G = \mathbb{Z}$. Under mild assumptions, we present a corresponding characterization for a general locally compact group G. Our results are obtained in a more general setting when the translations only by a subset $S \subset G$ are considered.

1. Introduction

Let G be a locally compact group with identity e and a left Haar measure μ. Fix p, $1 \leqslant p < \infty$, and a weight ω, that is, a measurable strictly positive function on G supposed to be locally p-summable.

Consider the weighted space $L^p(G,\omega) = \{f : \int_G |f\omega|^p d\mu < \infty\}$ of complex-valued functions on G. In this paper the following question is addressed. Can *all* functions in $L^p(G,\omega)$ be approximated arbitrarily well by the left translates of some *single* function $f \in L^p(G,\omega)$? The same question can be asked if we allow to translate f only by elements of a fixed subset $S \subset G$.

To be more precise, let S be a subset of G, and suppose that, for any $s \in S$, the (left) translation operator

$$(T_s f)(t) = f(s^{-1}t),\ t \in G$$

is continuous from $L^p(G,\omega)$ into itself. A function $f \in L^p(G,\omega)$ is called *S-dense* if its S-orbit $Orb_S(f) = \{T_s f : s \in S\}$ is dense in $L^p(G,\omega)$. So, we are interested in the existence of S-dense vectors in the space $L^p(G,\omega)$.

S-density is a particular case of a more general situation, known as the universality phenomenon, see a survey of K.-G. Grosse-Erdmann [8]. One of the first examples in this area was given in 1929 by G. D. Birkhoff (see [8]): there exists an entire function f whose translates are dense in the space $\mathcal{H}(\mathbb{C})$ of entire functions. In the case when S is the semigroup generated by one element $s \in G$, S-density is equivalent to the *hypercyclicity* of the operator T_s, which means that there exists a single vector x such that the sequence $(T_s^n x)_{n\in\mathbb{N}}$ is dense.

If S is a sub-semigroup or a subgroup of G, S-density means in other terms that the action of S by translations on the space $L^p(G,\omega)$ admits a hypercyclic vector. On a non-weighted L^p-space, the translation operators are isometric and therefore cannot have hypercyclic vectors. In the weighted case hypercyclicity can occur, depending on the weight; moreover, the conditions on weight are as a rule explicit and not difficult to calculate. This problem was considered by many authors. The characterisation in the discrete case $G = \mathbb{Z}$, $S = \mathbb{Z}_+$ was obtained by H. Salas in 1995 [10]. In 1997, W. Desch, W. Schappacher, and G. Webb [5] characterized the hypercyclicity of the C_0-semigroup of translations by $S = \mathbb{R}_+$, with $G = \mathbb{R}$. The dynamics of the translation C_0–semigroup indexed by a sector of the complex plane was considered by J. A. Conejero and A. Peris [2]. Ch.-Ch. Chen [1] characterized the hypercyclicity of a single translation operator (when S is the semigroup generated by an element of G) for locally compact groups.

In the present paper, we do not suppose that the set S has any algebraic structure: it can be an arbitrary subset of G. We denote by $\mathbf{\Sigma}$ the class of non-compact second countable locally compact

2010 *Mathematics Subject Classification.* 47A16; 37C85; 43A15.

Key words and phrases. locally compact groups; weighted spaces; hypercyclicity; translation semigroups.

groups. For a locally compact group G, we show that $G \in \mathbf{\Sigma}$ if and only if for some weight ω on G and for some $S \subset G$ there exist S-dense functions.

The main result of the paper is Theorem 5, which provides a necessary and sufficient condition for the existence of S-dense functions in $L^p(G,\omega)$. To avoid technical details, we formulate here this condition in the case when G is discrete:

Corollary 6. *Let G be an infinite countable discrete group, and let $S \subset G$. Let ω be a weight on G. There is an S-dense vector in $L^p(G,\omega)$ if and only if for every increasing sequence $(F_n)_{n\geqslant 1}$ of finite subsets of G, there exists a sequence $(s_n)_{n\geqslant 1} \subset S$ such that the sets $s_n^{-1}F_n$ are pairwise disjoint and, setting $s_0 = e$, $F_0 = \emptyset$,*

$$\sum_{n,k:n\neq k} \Big(\int_{s_n s_k^{-1} F_k} \omega^p \Big)^{1/p} < \infty.$$

If we suppose that S generates a commutative subgroup of G (which is not necessarily discrete), the criterion is straightforward:

Theorem 10. *Let $G \in \mathbf{\Sigma}$, and let $S \subset G$ generate an abelian subgroup in G. Let ω be a weight on G. There is an S-dense vector in $L^p(G,\omega)$ if and only if for every compact set $F \subset G$ and any given $\delta > 0$, there exist $s \in S$ and a compact $E \subset F$ such that $\mu(F \setminus E) < \delta$ and*

$$\operatorname*{ess\,sup}_{sE \cup s^{-1}E} \omega < \delta.$$

Note that if, in addition, G is discrete, compact subsets of G are just finite, and $E = F$ in this condition.

Our theorems generalize the results [5], [10], [1] mentioned above.

2. Notations and assumptions on the group

Recall that we consider the weighted space $L^p(G,\omega) = \{f : f\omega \in L^p(G)\}$ with the norm $\|f\|_{p,\omega} = \big(\int |f\omega|^p \big)^{1/p}$. The assumption that ω is locally p-summable (that is $\omega \in L^p(K)$ for any compact set $K \subset G$) implies that $L^p(G,\omega)$ contains all measurable compactly supported bounded functions. For a set $K \subset G$, denote by I_K the characteristic function of K. For a measurable function f, denote $\|f\|_{p,K} = (\int_K |f|^p)^{1/p}$, with $\|f\|_p = \|f\|_{p,G}$.

The Haar measure is denoted by μ, but we will also use the notations $|A| \equiv \mu(A)$ for $A \subset G$ and $\int_G f d\mu \equiv \int_G f(t)\, dt$. The translation $T_s f$ of a function $f \in L^p(G,\omega)$ will also be denoted by ${}^s f$ ($s \in G$).

For $s \in G$, one has the following expression for the norm of the translation operator:

$$\|T_s\| = \operatorname*{ess\,sup}_{t\in G} \frac{\omega(st)}{\omega(t)}. \tag{1}$$

We say that a weight ω is *S-admissible* if $\|T_s\| < +\infty$ for all $s \in S$.

Let $\mathbf{\Sigma}$ be the class of non-compact second countable locally compact groups. Note that every group in $\mathbf{\Sigma}$ is σ-compact, that is, represented as a countable union of compact sets (to see this, pick an open compactly generated subgroup $H \subset G$, which is a fortiori σ-compact; G being second countable, $[G : H]$ is necessarily countable, so finally G is σ-compact). In this section we show that G must be in $\mathbf{\Sigma}$ to admit S-dense vectors for some $S \subset G$ and some weight ω.

In the statements 1-3 it is supposed that $\S \subset G$ and the weight ω is S-admissible.

Our first claim is that G can't be compact. In the following lemma we prove a bit more, and it is exactly this result that we use below in the proof of Theorem 5.

Lemma 1. *Let x be an S-dense vector in $L^p(G,\omega)$ for some S, p, ω, and let $K \subset G$ be a compact set. Then for every $\varepsilon > 0, \lambda \in \mathbb{R}$ and every compact set $L \subset G$ there is $s \in S \setminus L$ such that $\|\lambda I_K - {}^s x\|_{p,\omega} < \varepsilon$.*

Proof. We can scale the Haar measure if necessary to have $\mu(K) = 1$. Increasing L if necessary, we can suppose that $L = L^{-1}$. Suppose the contrary: that is, $\|\lambda I_K - {}^s x\|_{p,\omega} \geqslant \varepsilon$ for every $s \in S \setminus L$. There is clearly $C > 0$ such that for $K_C = \{t \in K : \omega(t) > 1/C\}$ we have $\mu(K_C) > 3/4$.

Since LK is also compact, its measure is finite. Pick $N \in \mathbb{N}$ with $N > 2\mu(LK)$. Set $\lambda_j = \lambda + j\,\varepsilon/(2N)$, $j = 0, \dots, N$ and pick $\delta > 0$ such that $\delta < \varepsilon^p/(4N)^p$ and $\delta/C^p < \varepsilon$. For $j = 0, \dots, N-1$,

there is $s_j \in S$ such that $\|\lambda_j I_K - {}^{s_j}x\|_{p,\omega}^p < \delta/(4C^p)$. Since $\|\lambda I_K - {}^{s_j}x\|_{p,\omega} < \delta/(4C^p) + |\lambda - \lambda_j| < \varepsilon/4 + \varepsilon/2 < \varepsilon$, by assumption we have $s_j \in L$.

It follows that

$$\int_{s_j K_C} |\lambda_j - x(t)|^p dt = \int_{K_C} |\lambda_j - x(s_j^{-1}t)|^p dt < C^p \int_{K_C} \omega(t)^p |\lambda_j - x(s_j^{-1}t)|^p dt$$
$$\leqslant C^p \|\lambda_j I_K - {}^{s_j}x\|_{p,\omega}^p < C^p \frac{\delta}{4C^p} = \frac{\delta}{4}.$$

Set $K_j = \{t \in K_C : |\lambda_j - x(s_j^{-1}t)|^p \geqslant \delta\}$. By trivial estimates,

$$\frac{\delta}{4} > \int_{K_j} |\lambda_j - x(s_j^{-1}t)|^p dt \geqslant \delta\, \mu(K_j),$$

so that

$$\mu(K_j) < \frac{1}{4}.$$

For $t \in K_j' = s_j^{-1}(K_C \setminus K_j)$ we have the estimate $|x(t) - \lambda_j| < \delta^{1/p} < \varepsilon/(4N)$. As $|\lambda_j - \lambda_k| \geqslant \varepsilon/(2N)$ for $j \neq k$, it follows that the sets K_j' are pairwise disjoint. Moreover, $\mu(K_j') \geqslant \mu(K_C) - \mu(K_j) > 1/2$ for every j, hence $\mu(\cup K_j') > N/2 > \mu(LK)$ and in particular, $\cup K_j' \not\subset LK$. But by assumption, we have $K_j' \subset s_j^{-1}K \subset LK$ for every j. This contradiction proves the lemma. □

Corollary 2. *If for some p, S, ω there exists an S-dense vector in $L^p(G,\omega)$, then G is not compact.*

Proposition 3. *If for some p, S, ω there exists an S-dense vector in $L^p(G,\omega)$, then G is second countable.*

Proof. We use then the fact that G is second countable if and only if $L^p(G)$ is separable [4, Theorem 2], and the same is valid for $L^p(G,\omega)$ since this space is isometrically isomorphic to $L^p(G)$.

Suppose that $L^p(G,\omega)$ is not separable but $x \in L^p(G,\omega)$ is an S-dense vector. There exists then an uncountable set $\{f_\alpha\} \subset L^p(G,\omega)$ such that $\|f_\alpha - f_\beta\|_p > 3$ (this is easy to show usign the Zorn's lemma). Approximating f_α by ${}^{s_\alpha}x$ so that $\|f_\alpha - {}^{s_\alpha}x\|_p < 1/2$, we get a family $s_\alpha \in S$ such that $\|{}^{s_\alpha}x - {}^{s_\beta}x\|_{p,\omega}^p > 2$. In particular, s_α are all different.

For every α,

$$\|{}^{s_\alpha}x\|_{p,\omega}^p = \int \omega^p |{}^{s_\alpha}x|^p = \lim_{N\to\infty} \int_{\omega\leqslant N} \omega^p |{}^{s_\alpha}x|^p.$$

There exists $N_\alpha \in \mathbb{N}$ such that $\int_{\omega\geqslant N_\alpha} \omega^p |{}^{s_\alpha}x|^p < 1/2$.

Since the set of α is uncountable, there is a single $N \in \mathbb{N}$ such that the set $A = \{\alpha : N_\alpha = N\}$ is uncountable. For every $\alpha, \beta \in A$, we have

$$\int_{\omega\leqslant N} |{}^{s_\alpha}x - {}^{s_\beta}x|^p \geqslant N^{-p} \int_{\omega\leqslant N} \omega^p |{}^{s_\alpha}x - {}^{s_\beta}x|^p$$
$$\geqslant N^{-p}\Big(\|{}^{s_\alpha}x - {}^{s_\beta}x\|_{p,\omega}^p - \int_{\omega\geqslant N_\alpha} \omega^p |{}^{s_\alpha}x|^p - \int_{\omega\geqslant N_\alpha} \omega^p |{}^{s_\beta}x|^p\Big) > N^{-p}.$$

But this contradicts the fact that the G-orbit of any function in the non-weighted $L^p(G)$ is separable [11]: pick $\gamma \in A$ and set $y = {}^{s_\gamma}x \cdot I_{\omega\geqslant N}$, then $y \in L^p(G)$ (with norm $\leqslant N/2^{1/p}$), and $\int_{\omega\leqslant N} |{}^{s_\alpha}x - {}^{s_\beta}x|^p = \|{}^{(s_\alpha s_\gamma^{-1})}y - {}^{(s_\beta s_\gamma^{-1})}y\|_p^p$ for all $\alpha \neq \beta$ in A. □

Corollary 2 and Proposition 3 prove the implication (2) ⇒ (1) of the following

Theorem 4. *For a locally compact group G, the following assumptions are equivalent:*

1. *G is in the class $\mathbf{\Sigma}$ (non-compact second countable);*
2. *for every subset $S \subset G$ with non-compact closure there is an S-admissible weight ω such that $L^p(G,\omega)$ contains S-dense vectors, for all (or some) $p \geqslant 1$;*
3. *there is a weight ω for which all left and right translations are bounded and such that $L^p(G,\omega)$ contains G-dense functions for all (or some) $p \geqslant 1$.*

The implication (1) ⇒ (3) is proved in the beginning of Section 5 (and (3) ⇒ (2) is obvious).

3. Density criterion

In the most general case, the criterion has the following form. Below we give similar conditions in several particular cases. Recall that in every statement we assume that T_s is a bounded operator on $L^p(G,\omega)$ for all $s \in S$.

Theorem 5. *Let $G \in \mathbf{\Sigma}$, and let $S \subset G$. Let ω be an S-admissible weight on G. There is an S-dense vector in $L^p(G,\omega)$ if and only if for every increasing sequence $(F_n)_{n\geqslant 1}$ of compact subsets of G and any given $\delta_n > 0$, there exists a sequence $(s_n)_{n\geqslant 1} \subset S$ and compact sets $K_n \subset F_n$ such that the sets $s_n^{-1}F_n$ are pairwise disjoint, $|F_n \setminus K_n| < \delta_n$ and, setting $s_0 = e$, $K_0 = \emptyset$,*

$$\sum_{n,k\geqslant 0: n\neq k} \|\omega\|^p_{p, s_n s_k^{-1} K_k} < \infty. \tag{2}$$

Proof. $(\Rightarrow)$:
We set $s_0 = e$ and $K_0 = \emptyset$.

Decreasing δ_k if necessary, we can assume that $\sum \delta_k < \infty$. Let $y \in L^p(G,\omega)$ be S-dense. For every k, there is $F_k' \subset F_k$ such that $|F_k \setminus F_k'| < \delta_k/2$ and

$$\operatorname*{ess\,inf}_{F_k'} \omega \geqslant c_k, \qquad \operatorname*{ess\,sup}_{F_k'} \omega \leqslant c_k^{-1}$$

with some $0 < c_k < 1/4$. By Lemma 1, there exists $s_k \in S$ such that $\|2I_{F_k'} - {}^{s_k}y\|^p_{p,\omega} < c_k^{2p}\,\delta_k$ and $s_k^{-1}F_k$ does not intersect $s_j^{-1}F_j$ for $j < k$ (this condition is equivalent to $s_k \notin F_k F_n^{-1} s_n$). This implies that

$$c_k^{2p}\,\delta_k > \int_{F_k'} |{}^{s_k}y(t) - 2|^p\,\omega(t)^p dt \geqslant c_k^p \int_{F_k'} |{}^{s_k}y(t) - 2|^p dt$$

and so $\|{}^{s_k}y - 2\|^p_{p,F_k'} < c_k^p\delta_k < \delta_k$.

Set $K_k' = \{t \in F_k' : |y(s_k^{-1}t)| > 1\}$. Then

$$c_k^p\delta_k > \|{}^{s_k}y - 2\|^p_{p,F_k'} = \int_{K_k'} |y(s_k^{-1}t) - 2|^p dt + \int_{F_k'\setminus K_k'} |y(s_k^{-1}t) - 2|^p dt, \tag{3}$$

which implies

$$|F_k' \setminus K_k'| \leqslant \int_{F_k'\setminus K_k'} |y(s_k^{-1}t) - 2|^p dt < c_k^p\delta_k < \delta_k/4.$$

By regularity of the Haar measure, there is a compact set $K_k \subset K_k'$ such that $|K_k' \setminus K_k| < \delta_k/4$. By estimates above, we have $|F_k \setminus K_k| < \delta_k$.

Since the norm of y is finite, we have:

$$\sum_k \int_{s_k^{-1}K_k} \omega^p \leqslant \sum_k \int_{s_k^{-1}K_k} \omega^p(t)\,|y(t)|^p dt \leqslant \|y\|^p_{p,\omega} < \infty.$$

By the choice of s_n, we have

$$\begin{aligned}\int_{G\setminus K_n} \omega^p\,|{}^{s_n}y|^p &= \int_{F_n'\setminus K_n} \omega^p\,|{}^{s_n}y|^p + \int_{G\setminus F_n'} \omega^p\,|{}^{s_n}y|^p \\ &\leqslant |F_n' \setminus K_n| c_n^{-p} + \|2I_{F_n'} - {}^{s_n}y\|^p_{p,\omega} < \delta_n + c_n^p\,\delta_n \leqslant 2\delta_n.\end{aligned}$$

At the same time, $|{}^{s_n}y|^p > 1$ on $s_n s_k^{-1}K_k$. For $k \neq n$, the condition $s_n s_k^{-1}F_k \cap F_n = \emptyset$ implies that $s_n s_k^{-1}K_k \subset G \setminus K_n$; moreover, the sets $s_n s_k^{-1}F_k$, $k \neq n$, are pairwise disjoint. It follows that

$$2\delta_n > \sum_{k:k\neq n} \int_{s_n s_k^{-1}K_k} \omega^p\,|{}^{s_n}y|^p > \sum_{k:k\neq n} \|\omega\|^p_{p, s_n s_k^{-1}K_k}.$$

By the choice of δ_n we have $\sum_n \delta_n < \infty$, so we get immediately (2).

$(\Leftarrow)$:
From the assumptions it follows that $L^p(G,\omega)$ is separable. Choose a dense sequence $(Q_n)_{n=1}^\infty \subset L^p(G,\omega)$ such that every Q_n is compactly supported and essentially bounded. Arrange them with repetitions in a sequence $(P_n)_{n=1}^\infty$ so that every Q_n appears infinitely many times in this sequence.

We will seek for X "almost" of the form $\sum_k T_{s_k^{-1}} P_k$, with $s_n \in S$ chosen so that the series converges and $\|^{s_n} X - P_n\|_{p,\omega} < \varepsilon_n$ for every n, with some $\varepsilon_n \to 0$. This guarantees that X is S-dense.

Denote $F_n = \operatorname{supp} P_n$ and set $p_n = \|P_n\|_\infty^p$. Pick a decreasing sequence (δ_n) such that $0 < \delta_n < p_n^{-1} 2^{-n}$. Choose s_n, K_n according to (2). Set

$$a_k = \sum_{n:n\neq k} \|\omega\|^p_{p,s_n s_k^{-1} K_k}.$$

By (2), $\sum_k a_k < \infty$. Choose a subsequence (a_{l_k}), with $l_0 = 0$, such that $\sum p_k a_{l_k} < \infty$ and $P_{l_k} = P_k$ for every k. By the choice of (P_k), this is always possible. Set $t_n = s_{l_n}$ and $E_k = K_{l_k}$, then the following series converges:

$$\sum_{n,k:n\neq k} p_k \|\omega\|^p_{p,t_n t_k^{-1} E_k} < \infty. \tag{4}$$

In particular, the sets $t_n^{-1} E_n$ are pairwise disjoint, and

$$|\operatorname{supp} P_n \setminus E_n| = |\operatorname{supp} P_{l_n} \setminus K_{l_n}| < \delta_{l_n} \leqslant \delta_n$$

for $n \geqslant 1$. We can still assume that $t_0 = e$, $E_0 = \emptyset$. Set

$$X = \sum_k T_{t_k^{-1}}(P_k I_{E_k}).$$

This series converges, since (note that the support of $T_{t_k^{-1}} P_k$ is $t_k^{-1} \operatorname{supp} P_k$)

$$\begin{aligned}\|X\|^p_{p,\omega} &= \sum_k \int |P_k(t_k t) I_{E_k}(t_k t)\omega(t)|^p dt \leqslant \sum_k \|P_k\|^p_\infty \int_{t_k^{-1}E_k} |\omega(t)|^p dt \\ &= \sum_k p_k \|\omega\|^p_{p,t_0 t_k^{-1} E_k} < \infty.\end{aligned}$$

Now, for every n

$$\|^{t_n} X - P_n\|^p_{p,\omega} = \|P_n(1 - I_{E_n})\|^p_{p,\omega} + \sum_{k:k\neq n} \|^{t_n t_k^{-1}}(P_k I_{E_k})\|^p_{p,\omega}, \tag{5}$$

since again the supports of the summands in (5) are pairwise disjoint:

$$\operatorname{supp}{}^{t_n t_k^{-1}} P_k \subset t_n t_k^{-1} E_k.$$

One has

$$\|P_n(1 - I_{E_n})\|^p_{p,\omega} = \int_{\operatorname{supp} P_n \setminus E_n} |P_n|^p \leqslant \|P_n\|^p_\infty |\operatorname{supp} P_n \setminus E_n| < p_n \delta_n < 2^{-n}.$$

and for $k \neq n$,

$$\|^{t_n t_k^{-1}}(P_k I_{E_k})\|^p_{p,\omega} \leqslant \|P_k\|^p_\infty \|\omega\|^p_{p,t_n t_k^{-1} E_k} = p_k \|\omega\|^p_{p,t_n t_k^{-1} E_k}.$$

Denote

$$\varepsilon_n = \sum_{n,k:k\neq n} p_k \|\omega\|^p_{p,t_n t_k^{-1} E_k}.$$

By (4), $\sum_n \varepsilon_n < \infty$, so $\varepsilon_n \to 0$. Thus, we have

$$\|^{t_n} X - P_n\|^p_{p,\omega} \leqslant \varepsilon_n + 2^{-n} \tag{6}$$

with $\varepsilon_n \to 0$, which proves that X is S-dense. □

Corollary 6. *Let G be an infinite countable discrete group, and let $S \subset G$. Let ω be an S-admissible weight on G. There is an S-dense vector in $L^p(G,\omega)$ if and only if for every increasing sequence $(F_n)_{n\geqslant 1}$ of finite subsets of G, there exists a sequence $(s_n)_{n\geqslant 1} \subset S$ such that the sets $s_n^{-1} F_n$ are pairwise disjoint and, setting $s_0 = e$, $F_0 = \emptyset$,*

$$\sum_{n,k\geqslant 0:n\neq k} \|\omega\|^p_{p,s_n s_k^{-1} F_k} < \infty. \tag{7}$$

Remark 7. *It is enough to check the conditions of Theorem 5 for an increasing sequence (F_n) of compact sets such that $G = \cup_n F_n$. Indeed, for another sequence $(\tilde{F}_k)$ one can choose n_k so that $\tilde{F}_k \subset F_{n_k}$, and the rest is obvious.*

4. Simplifications of conditions

Theorem 8. *Let $G \in \mathbf{\Sigma}$, and let $S \subset G$. Let ω be an S-admissible weight on G. If for every compact set $K \subset G$ of positive measure and every $\varepsilon, \delta > 0$ there exist $s \in S$ and compact $E \subset K$ such that $|K \setminus E| < \delta$ and*

$$\operatorname*{ess\,sup}_{sE \cup s^{-1}E} \omega < \varepsilon, \tag{8}$$

then $L^p(G, \omega)$ contains an S-dense vector.

Proof. Let us show first that under the assumptions above, s can be chosen outside of a given compact set L. Suppose the contrary: that (8) does not hold for any $s \notin L$. We can assume that $\delta_1 = |K| - \delta > 0$. Choose a compact set E_0 and $s_0 \in S$ so that $|K \setminus E_0| < \delta$ and (8) holds for $E = E_0$, $s = s_0$. In particular, $|E_0| > |K| - \delta = \delta_1$. There is a compact subset $K_1 \subset E_0$ such that $|E_0 \setminus K_1| < \delta_1/9$ and $\varepsilon_1 = \operatorname{ess\,inf}\{\omega(t) : t \in s_0 K_1 \cup s_0^{-1} K_1\} > 0$.

Next choose by induction compact sets E_n, K_n and $s_n \in S$, $n \geqslant 1$, so that $|K_n \setminus E_n| < \delta_1/9^n$, $\operatorname{ess\,sup}_{s_n E_n \cup s_n^{-1} E_n} \omega < \varepsilon_n$, $|E_n \setminus K_{n+1}| < \delta_1/9^n$ and $\varepsilon_{n+1} = \operatorname{ess\,inf}\{\omega(t) : t \in s_n K_{n+1} \cup s_n^{-1} K_{n+1}\} > 0$. For every $n \geqslant 1$, we have $|E_n| > |K_n| - \delta_1/9^n > |E_{n-1}| - 2\delta_1/9^n$, which implies

$$|E_n| > |E_0| - 2\delta_1 \sum_{k=1}^{n} \frac{1}{9^k} > |E_0| - \delta_1/4 = 3\delta_1/4.$$

Denote $D_n = s_n E_n \cup s_n^{-1} E_n$, $D_n^- = \cup_{k<n} D_k \cap D_n$ and $D_n^+ = \cup_{k>n} D_k \cap D_n$. Since $D_k \cap D_n$ is contained (except probably for a zero measure subset) in $(\{s_n\} \cup \{s_n^{-1}\})(E_n \setminus K_{n+1})$ for all $k > n$, we have $|D_n^+| \leqslant 2|E_n \setminus K_{n+1}| < 2\delta_1/9^n$. As a consequence,

$$|D_n^-| \leqslant \sum_{k<n} |D_k^+| < 2\delta_1 \sum_{k=1}^{\infty} \frac{1}{9^k} = \frac{\delta_1}{8},$$

and for every n,

$$\begin{aligned} |\cup_{k \leqslant n} D_k| &= |\cup_{k<n} D_k \sqcup (D_n \setminus D_n^-)| \\ &\geqslant |\cup_{k<n} D_k| + |D_n| - |D_n^-| \geqslant |\cup_{k<n} D_k| + \frac{3\delta_1}{4} - \frac{\delta_1}{8} > |\cup_{k<n} D_k| + \frac{\delta_1}{2}, \end{aligned}$$

whence it follows that

$$|LE_0 \cup L^{-1}E_0| \geqslant |\cup_{k=1}^{\infty} D_k| \geqslant \sum_{k=1}^{\infty} \frac{\delta_1}{2} = +\infty,$$

which is impossible since $LE \cup L^{-1}E$ is compact. This contradiction shows that s can be chosen outside L.

Now let us show that the assumptions of the theorem imply the assumptions of Theorem 5.

Let (F_n) be an increasing sequence of compact sets and (δ_n) a sequence of positive numbers. We can assume that $\delta_1 < 1/2$ and $\delta_{n+1} < \delta_n/2$ for all n, in particular $\delta_n < 2^{-n}$. Choose firstly sequences $E_n' \subset G$, $s_n \in S$ by induction. Set $E_0' = \emptyset$, $s_0 = e$; for $n \geqslant 1$, set

$$K_n = \Big(\bigcup_{k<n} s_k F_n \Big) \cup \Big(\bigcup_{k<n} F_k \cup s_k^{-1} F_k \Big).$$

Set $C_n = \max_{k<n} \|T_{s_k}\|$ (automatically $C_n \geqslant 1$) and choose $s_n \in S$, $E_n' \subset K_n$ so that E_n' is compact, $|K_n \setminus E_n'| < \delta_n/2$ and $\|\omega\|_{\infty, s_n E_n' \cup s_n^{-1} E_n'} < (2^n C_n |E_n'|^{1/p})^{-1}$. We can choose s_n so that $s_n^{-1} F_n$ does not intersect $s_k^{-1} F_k$ for $k < n$ (that is, $s_n \notin F_n F_k^{-1} s_k$).

Set now $E_0 = \emptyset$ and

$$E_n = F_n \cap E_n' \cap s_n(\cap_{k>n} E_k') \tag{9}$$

for $n \geqslant 1$. Then, since $F_n \subset K_n$ and for $k > n$ $s_n^{-1}F_n \subset K_k$, that is $F_n \subset s_n K_k$, we have

$$|F_n \setminus E_n| \leqslant |K_n \setminus E_n'| + |\cup_{k>n} s_n K_k \setminus s_n E_k'| = \sum_{k \geqslant n} |K_k \setminus E_k'|$$
$$< \sum_{k \geqslant n} \frac{\delta_k}{2} < \sum_{k \geqslant n} \frac{\delta_n}{2^{k-n+1}} = \delta_n.$$

The sets $s_n^{-1}F_n$ are pairwise disjoint; for $k < n$ we have $E_k \subset s_k E_n'$ by (9), which implies $s_k^{-1}E_k \subset s_k^{-1}F_k \subset E_n'$. Now

$$\begin{aligned}\sum_{n,k:n\neq k} \|\omega\|^p_{p,s_n s_k^{-1}E_k} &= \sum_n \left(\sum_{k<n} \|\omega\|^p_{p,s_n s_k^{-1}E_k} + \sum_{k>n} \|\omega\|^p_{p,s_n s_k^{-1}E_k} \right) \\ &\leqslant \sum_n \left(\|\omega\|^p_{p,s_n E_n'} + \sum_{k>n} \|T_{s_n}\|^p \|\omega\|^p_{p,s_k^{-1}E_k} \right) \\ &\leqslant \sum_n \left(\|\omega\|^p_{\infty,s_n E_n'} |E_n'| + \sum_{k>n} C_k^p \|\omega\|^p_{\infty,s_k^{-1}E_k'} |E_k'| \right) \\ &= \sum_n \left(2^{-n} + \sum_{k>n} 2^{-k} \right) < \infty.\end{aligned}$$

Finally, the assumptions of Theorem 5 are satisfied with E_n in place of K_n. □

Corollary 9. *Let $G \in \mathbf{\Sigma}$, and let $S \subset G$. Let ω be an S-admissible weight on G. If for every compact set $K \subset G$ of positive measure*

$$\inf_{s \in S} \left(\operatorname*{ess\,sup}_{sK \cup s^{-1}K} \omega \right) = 0, \tag{10}$$

then $L^p(G,\omega)$ contains an S-dense vector.

In general, this condition is not necessary, see Example 15. But if S generates an abelian subgroup, this is the case:

Theorem 10. *Suppose that $G \in \mathbf{\Sigma}$, and let $S \subset G$ generate an abelian subgroup. Let ω be an S-admissible weight on G. The following conditions are equivalent:*

1. *There is an S-dense vector in $L^p(G,\omega)$ for every p, $1 \leqslant p < \infty$;*
2. *There is an S-dense vector in $L^p(G,\omega)$ for some p, $1 \leqslant p < \infty$;*
3. *For every compact set $F \subset G$ and any given $\delta > 0$, there exist $s \in S$ and a compact $E \subset F$ such that $|F \setminus E| < \delta$ and*

$$\operatorname*{ess\,sup}_{sE \cup s^{-1}E} \omega < \delta; \tag{11}$$

4. *For some p, $1 \leqslant p < \infty$, the following condition holds: For every compact set $F \subset G$ and any given $\delta > 0$, there exist $s \in S$ and a compact $E \subset F$ such that $|F \setminus E| < \delta$ and*

$$\|\omega\|_{p,sE \cup s^{-1}E} < \delta. \tag{12}$$

5. *For every $1 \leqslant p < \infty$, the assumptions of (4) above hold.*

Proof. (4)⇒(3) is an easy exercise, (3)⇒(1) was shown in Theorem 8 without the commutativity assumption; (1)⇒(2) and (5) ⇒ (4) are obvious.

It remains to prove (2)⇒(5).

If there is an S-dense vector in $L^p(G,\omega)$, then for $F_n = F$ and $\delta_n = 2^{-n}$, $n \geqslant 1$, there exist $s_n \in S$, $E_n \subset F$ with $s_0 = e$, $E_0 = \emptyset$ such that $|F \setminus E_n| < \delta_n$ and

$$\sum_{n,k:n\neq k} \|\omega\|^p_{p,s_n s_k^{-1}E_k} < \infty.$$

It follows (by considering the subseries with $n = 0$) that $\|\omega\|^p_{p,s_k^{-1}E_k} \to 0$.

Let k be such that $\delta_k < \delta/2$. Set $C = \|T_{s_k}\|$. For $n > k$, we have

$$\|\omega\|^p_{p,s_nE_k} = \int_{s_nE_k} |\omega|^p = \int_{s_ks_ns_k^{-1}E_k} |\omega|^p \leqslant C \int_{s_ns_k^{-1}E_k} |\omega|^p \to 0, \quad n \to \infty.$$

Pick now $n > k$ such that $\|\omega\|_{p,s_n^{-1}E_n} < \delta/2$ and $\|\omega\|_{p,s_nE_k} < \delta/2$. Set $E = E_k \cap E_n$, $s = s_n$, then $|F \setminus E| < \delta_k + \delta_n < \delta$ and $\|\omega\|_{p,sE\cup s^{-1}E} \leqslant \|\omega\|^p_{p,s_nE_k} + \|\omega\|^p_{p,s_n^{-1}E_n} < \delta$. □

The following case is quite particular and is meaningful especially in the abelian case, see Corollary 13.

Remark 11. *If ω is locally summable and*

$$L(g) = \sup_{t\in G} \frac{\omega(gt)}{\omega(t)} < +\infty \tag{13}$$

for all $g \in G$, then ω is equivalent to a continuous weight [7, Satz 2.7]. *Since* (13) *holds or does not hold simlutaneously for ω^p for all $1 \leqslant p < \infty$, we have the same conclusion if ω is locally p-summable and satisfies* (13). *Hence in the following statement we can assume, without loss of generality, that ω is continuous.*

Proposition 12. *Let $G \in \mathbf{\Sigma}$, and suppose that the weight ω is continuous and such that for all $g \in G$,*

$$L(g) = \|T_g\| = \sup_{t\in G} \frac{\omega(gt)}{\omega(t)} < +\infty, \qquad R(g) = \sup_{t\in G} \frac{\omega(tg)}{\omega(t)} < +\infty. \tag{14}$$

Then for any $S \subset G$, the following conditions are equivalent:

(1) *There are S-dense vectors in $L^p(G,\omega)$.*

(2)

$$\inf_{s\in S} \sup_{sK\cup s^{-1}K} \omega = 0 \quad \textit{for every compact set } K \subset G \tag{15}$$

(3)

$$\inf_{s\in S} \max\big(\omega(s), \omega(s^{-1})\big) = 0 \tag{16}$$

Proof. Obviously (15)⇒(16). Let us prove the inverse implication. Let K be a compact set; we can assume that $K = K^{-1}$. It is proved in a rather general situation in [6, Proposition 1.16] that L and R are locally bounded (provided they are finite). There exists thus a constant C such that $L|_K \leqslant C$ and $R|_K \leqslant C$.

Now for every $s \in G$,

$$\sup_{t\in sK} \omega(t) \leqslant \sup_{t\in K} R(t)\omega(s) \leqslant C\omega(s).$$

In the same way we get the inequality $\sup_{s^{-1}K} \omega \leqslant C\omega(s^{-1})$, thus (15) and (16) are equivalent.

Since they both imply (10), they imply also the assumption (2) of Theorem 5 and the existence of S-dense vectors in $L^p(G,\omega)$.

Suppose now that there exist S-dense vectors in $L^p(G,\omega)$. Pick a compact set F of positive measure and set $F_n = F$ for all $n \in \mathbb{N}$. Choose $s_n \in S$ and $K_n \subset F$ so that (2) holds. By setting $\delta_n = |F|/4$, we can guarantee that $|K_n \cap K_1| > |F|/2$ for all $n \geqslant 1$. From (2) it follows, in particular, that

$$\lim_{n\to\infty} \|\omega\|^p_{p,s_ns_1^{-1}K_1} = 0.$$

Let $C_1 > 0$ be such that $L(s_1^{-1}) \leqslant C_1$, $R|_{K_1^{-1}} \leqslant C_1$ and $R|_{K_1^{-1}s_1} \leqslant C_1$. We have

$$\omega(s_n) = \inf_{t\in s_1^{-1}K_1} \omega(s_ntt^{-1}) \leqslant \inf_{t\in s_1^{-1}K_1} R(t^{-1})\omega(s_nt) \leqslant C_1\|\omega\|_{p,s_ns_1^{-1}K_1}|K_1|^{-1}$$

which implies $\omega(s_n) \to 0$, $n \to \infty$. At the same time,

$$\begin{aligned}\omega(s_n^{-1}) &= \inf_{t\in K_1\cap K_n} \omega(s_1^{-1}s_1s_n^{-1}tt^{-1}) \leqslant \inf_{t\in K_1\cap K_n} L(s_1^{-1})\omega(s_1s_n^{-1}t)R(t^{-1}) \\ &\leqslant C_1^2 \inf_{t\in K_1\cap K_n} \omega(s_1s_n^{-1}t) \leqslant C_1^2\|\omega\|_{p,s_1s_n^{-1}K_n}|K_1\cap K_n|^{-1} < 2C_1^2|F|^{-1}\|\omega\|_{p,s_1s_n^{-1}K_n}.\end{aligned}$$

By (2), $\omega(s_n^{-1}) \to 0$, $n \to \infty$, and this implies (16). □

In the case when G is abelian our conditions take an especially simple form:

Corollary 13. *Let $G \in \mathbf{\Sigma}$ be abelian, $1 \leqslant p < \infty$, and let ω be a continuous G-admissible weight. For any given $S \subset G$, there are S-dense vectors in $L^p(G,\omega)$ if and only if*

$$\inf_{s\in S} \max\big(\omega(s), \omega(s^{-1})\big) = 0. \tag{17}$$

5. Examples and counter-examples

First we show that every group $G \in \mathbf{\Sigma}$ admits a weight with G-dense vectors; this completes the proof of Theorem 4.

Proof. Let $U = U^{-1}$ be a neighbourhood of identity with compact closure and let G_1 be the subgroup generated by U. Pick $g_n \in G$, $n \in \mathbb{N}$, such that $G = \cup_n (g_n G_1)$; this is possible since G is σ-compact. Set $U_1 = U$ and for $n > 1$ set by induction

$$U'_n = \{g_n\} \cup \big(\bigcup_{k=1}^{n-1} U_k U_{n-k} \big), \qquad U_n = U'_n \cup (U'_n)^{-1}.$$

In particular, we have $U^n \subset U_n$ and $G = \cup_n U_n$. By construction, $U_j U_k \subset U_{j+k}$ for all j, k.

Every U_n has compact closure, so we can choose $s_n \in S$, $n \in \mathbb{N}$, so that $E_n = U_{n+1} s_n U_{n+1} \cup U_{n+1} s_n^{-1} U_{n+1}$ are pairwise disjoint and $U_{n+1} s_n U_{n+1} \cap U_{n+1} s_n^{-1} U_{n+1} = \emptyset$. Denote $V_{n,k} = U_k s_n U_k \cup U_k s_n^{-1} U_k$ for $1 \leqslant k \leqslant n$, $V_{n,k} = G$ for $k > n$ and $V_{n,0} = \{s_n\} \cup \{s_n^{-1}\}$. We have $U_j \, (V_{n,k} \setminus V_{n,k-1}) U_j \subset V_{n,k+j} \setminus V_{n,k-j-1}$ for all n, k. Set

$$\omega(s) = \begin{cases} 1, & s \notin \bigcup_n V_{n,n}, \\ 2^{-n+k}, & s \in V_{n,k} \setminus V_{n,k-1} \quad (k \leqslant n). \end{cases}$$

We have thus $\omega(s_n) = \omega(s_n^{-1}) = 2^{-n}$ and $\omega(t) \leqslant 1$ for all $t \in G$.

Let t be in U_j. If $s \notin \cup_n V_{n,n}$, then $st \notin \cup_{n;\, k \leqslant n-j-1} V_{n,k}$ and $\omega(st) \geqslant 2^{-j}$. This implies

$$\frac{1}{2^j} \leqslant \frac{\omega(st)}{\omega(s)} \leqslant 1.$$

Similarly $2^{-j} \leqslant \omega(ts)/\omega(s) \leqslant 1$. If $s \in V_{n,k} \setminus V_{n,k-1}$ with $k \leqslant n$, then $\omega(s) = 2^{k-n}$ and $st \in V_{n,k+j} \setminus V_{n,k-j-1}$, thus

$$\frac{1}{2^j} = \frac{2^{k-j-n}}{2^{k-n}} \leqslant \frac{\omega(st)}{\omega(s)} \leqslant \frac{2^{k+j-n}}{2^{k-n}} = 2^j.$$

Similarly $2^{-j} \leqslant \omega(ts)/\omega(s) \leqslant 2^j$, and we see that these inequalities hold for all $s \in G$, $t \in U$.

It follows that in the notations of (14), $L(t) < +\infty$, $R(t) < +\infty$ for all $t \in U_j$, and since $G = \cup_j U_j$, this is true for all $t \in G$. By construction, the condition (16) holds, so we can conclude that $L^p(G,\omega)$ contains an S-dense vector. □

Let $\mathbb{F}_2$ be the free discrete group of two generators a and b.

Example 14. *There exists an $\mathbb{F}_2$-admissible weight on $\mathbb{F}_2$ which satisfies the condition* (16) *but does not admit $\mathbb{F}_2$-dense vectors.*

Set $U = \{e, a, b, a^{-1}, b^{-1}\}$, then $G = \cup_n U^n$. Set $n_k = 2^{2^k}$, $k = 1, 2, \ldots$. Define the weight ω as follows: $\omega(a^{\pm n_k}) = 2^{-k}$, $\omega(t) = 2^{j-k}$ if $t \in (U^j \setminus U^{j-1})\, a^{\pm n_k}$, $1 \leqslant j \leqslant k$, and 1 elsewhere. In particular, $\omega|_U = 1$.

For $\alpha \in \{a, b, a^{-1}, b^{-1}\}$, denote by F_α the set of reduced words in $\mathbb{F}_2$ which end by α. Note that $U^k a^{\pm n_k} \subset F_a \cup F_{a^{-1}}$ for all k (since every word in U^k has length at most $k < n_k$). It follows that $\omega \equiv 1$ on $F_b \cup F_{b^{-1}}$.

Obviously $\omega(sg) \leqslant 2\omega(g)$ for $s \in U$ and for all $g \in \mathbb{F}_2$. It follows that ω is of moderate growth in the sense of Edwards [6], that is, $\|T_g\| < \infty$ for all $g \in \mathbb{F}_2$. Also, $\inf_g \max(\omega(g), \omega(g^{-1})) \leqslant \inf_k \max(\omega(a^{n_k}), \omega(a^{-n_k})) = 0$.

Suppose now that there is a $\mathbb{F}_2$-dense vector $x \in L^p(G,\omega)$. For every $0 < \varepsilon < 1/2$, there exists $g \in \mathbb{F}_2$ such that $\|I_U - {}^g x\|_{p,\omega} < \varepsilon$. This implies that

$$|x(g^{-1}t) - 1| \omega(t) < \varepsilon$$

for $t = e, b$. It follows that $|x(t)| > 1/2$ for $t = g^{-1}$ and $t = g^{-1}b$.

By Lemma 1, for the same ε there is $h \in \mathbb{F}_2 \setminus (Ug \cup Ub^{-1}g)$ such that $\|^h x - I_U\|_{p,\omega} < \varepsilon$. This implies that $|x(h^{-1}t)|\,\omega(t) < \varepsilon$ for $t \notin U$. In particular, setting $t = hg^{-1}$ we get: $|x(h^{-1}hg^{-1})|\,\omega(hg^{-1}) = |x(g^{-1})|\,\omega(hg^{-1}) < \varepsilon$ which implies $\omega(hg^{-1}) < 2\varepsilon < 1$; setting $t = hg^{-1}b$, we infer that $\omega(hg^{-1}b) < 1$. By construction of ω, this implies that $hg^{-1}, hg^{-1}b \notin F_b \cup F_{b^{-1}}$, which is impossible. This contradiction proves that there are no $\mathbb{F}_2$-dense vectors.

Example 15. *There exists an S-admissible weight on $G = \mathbb{F}_2$ which admits S-dense vectors in $L^p(G, \omega)$ for some semigroup S but does not satisfy the sufficient condition* (10).

Set $U = \{e, a, b, a^{-1}, b^{-1}\}$, then $G = \cup_n U^n$. Set $n_k = 2^{2^k}$, $k = 1, 2, \dots$.

For $k \in \mathbb{N}$, set $s_k = a^{n_k} b$ and let S be the semigroup generated by $\{s_k : k \in \mathbb{N}\}$. Set $V_{l,k} = s_l s_k^{-1} U^k$. Every element of $V_{l,k}$, $l \neq k$, has the form $s_l s_k^{-1} t = a^{n_l - n_k} t$ with some $t \in U^k$; in the reduced form it equals $a^p r$ with $|p| \geqslant |n_l - n_k| - k > 0$ and $r \in U^{2k}$ not starting with $a^{\pm 1}$.

Let us show that the sets $SV_{l,k}$, $k \neq l$, are pairwise disjoint. For $s = a^{n_{j_1}} b \dots a^{n_{j_m}} b$, $s' = a^{n_{j'_1}} b \dots a^{n_{j'_{m'}}} b$ and $a^p r \in V_{l,k}$, $a^{p'} r' \in V_{l',k'}$ (the latter two reduced as above) the equality $s a^p r = s' a^{p'} r'$ would mean

$$a^{n_{j_1}} b \dots a^{n_{j_m}} b a^p r = a^{n_{j'_1}} b \dots a^{n_{j'_{m'}}} b a^{p'} r',$$

and clearly both sides are in their reduced form. We can suppose that $m \leqslant m'$. It follows that $j_i = j'_i$ for $i = 1, \dots, m$. The equality $p = n_{j_{m+1}}$ is in fact impossible: it would imply $p > 0$, then $l > k$; but in this case $p \geqslant n_l - n_k - k > n_l/2 > n_{l-1}$ and $p \leqslant n_l - n_k + k < n_l$, so p cannot be equal to n_ν for any $\nu \in \mathbb{N}$. We conclude that $m = m'$, $s = s'$ and $p = p'$.

As discussed above, $p = n_l - n_k + \xi$ with $|\xi| \leqslant k$, and $p' = n_{l'} - n_{k'} + \xi'$ with $|\xi'| \leqslant k'$. Either $p = p' > 0$ and then $k > l$, $k' > l'$, or $p = p' < 0$ and $k < l$, $k' < l'$. In the first case we can suppose that $l \geqslant l'$. We have $n_l = n_{l'} + n_k - n_{k'} + \xi' - \xi$; by the choice of (n_k), an equality is possible only if $l = l'$. But then $n_k = n_{k'} + \xi - \xi'$, and for the same reason it follows that $k = k'$. In the second case one arrives similarly at the same conclusion. Finally, $l = l'$ and $k = k'$, which shows that $V_{l,k}$ are pairwise disjoint indeed.

Set now

$$\omega(t) = \begin{cases} 8^{-l-k}, & t \in SV_{l,k}; \\ 1, & \text{otherwise} \end{cases}$$

By the reasoning above, ω is well defined. We have

$$\sum_{n,k:n\neq k} \|\omega\|^p_{p, s_n s_k^{-1} U^k} \leqslant \sum_{n,k:n\neq k} 8^{-n-k}(4^k + 1) < \infty,$$

and this implies that the assumptions of Theorem 5 hold.

It remains to show that $\|T_{s_n}\| < \infty$ for all n. If $t \notin \cup_{l,k} SV_{l,k}$, then $\omega(t) = 1$ and $\omega(s_n t)/\omega(t) \leqslant 1$. If $t = s s_l s_k^{-1} u$ with $s \in S$ and $u \in U^k$, then $\omega(s_n t) = \omega(s_n s s_l s_k^{-1} u) = \omega(t)$. It follows that $\|T_{s_n}\| \leqslant 1$.

Example 16. *In general, one cannot have $G_n = F_n$ in Theorem 5, so that*

$$\sum_{n,k:n\neq k} \|\omega\|^p_{p, s_n s_k^{-1} F_k} < \infty. \tag{18}$$

This is shown below: there exists a weight ω on $G = \mathbb{R}^2$ which is S-admissible and such that the condition (11) *is satisfied for $S = \mathbb{R} \times \{0\}$, i.e. $L^p(G, \omega)$ contains an S-dense vector, but the strengthened condition* (18) *does not hold.*

Denote $\ell_0 = 0$ and $\ell_n = \sum_{k=1}^n k = n(n+1)/2$ for $n \geqslant 1$. Define ω as follows: $\omega(-x, y) = \omega(x, y)$, and for $x \geqslant 0$

$$\omega(x, y) = \begin{cases} 2^{-n}, & x \in [\ell_n, \ell_{n+1}),\ y \notin (1 - 2^{1-n}, 1),\ n \in \mathbb{Z} \\ 2^n, & x \in [\ell_n, \ell_{n+1}),\ y \in [1 - 2^{-n}, 1),\ n \in \mathbb{Z} \\ 2^{n-2k}, & x \in [\ell_n + k, \ell_n + k + 1),\ y \in (1 - 2^{1-n}, 1 - 2^{-n}),\ n \in \mathbb{Z}, 0 \leqslant k < n \end{cases}$$

This is illustrated as follows:



For all $t \in [0,1)$,

$$\sup_{x,y\in\mathbb{R}^2} \frac{\omega(x+t,y)}{\omega(x,y)} = \sup_{n\in\mathbb{Z},y\in[0,1]} \frac{\omega(n+1,y)}{\omega(n,y)};$$

it is clear from the picture above that this ratio is always bounded by 4. It follows that $\|T_{(t,0)}\| = 4$ for all $t \in [0,1)$, so that $S = \mathbb{R}\times\{0\}$ acts continuously on $L^p(\mathbb{R}^2)$.

Setting $F_n = [n, n+1]\times[0,1]$, we have $\int_{F_n}\omega^p \geqslant 1$ for all n, so clearly the condition (18) does not hold.

To check (11), it is sufficient to consider $F = [-n,n]\times[-n,n]$. Choose $t\in\mathbb{N}$ so that

$$t \geqslant 2, \quad 2^{-t} < \delta, \quad n2^{3-t} < \delta.$$

Set $E = F\setminus([-n,n]\times(1-2^{2-t},1))$. Then $|F\setminus E| = n2^{3-t} < \delta$.

Set $s = (2\ell_t, 0)$. Since $\ell_t > t$,

$$s + E \subset [\ell_t, +\infty)\times(\mathbb{R}\setminus(1-2^{2-t},1))$$

and

$$-s + E \subset (-\infty, -\ell_t]\times(\mathbb{R}\setminus[1-2^{1-t},1)).$$

This implies

$$\sup_{(s+E)\cup(-s+E)} \omega^p \leqslant 2^{-t} < \delta,$$

thus (11) holds.

If S generates G and G is abelian, then such an example is impossible, see Proposition 12.

In [2], it was shown on an example that a semigroup indexed by a complex sector can be hypercyclic without any single operator being hypercyclic. We provide a simple example illustrating the same situation:

Example 17. *There is a $\mathbb{Z}^2$-admissible weight on $\mathbb{Z}^2$ with $\mathbb{Z}^2$-dense vectors such that no single operator $T_{(m,n)}$ is hypercyclic. The weight is defined as follows: set $\omega(l,m) = 2^{k-n}$ if $\max(|l-n|, |m-2^n|) = k$ or $\max(|l+n|, |m+2^n|) = k$ for some $n\in\mathbb{N}$ and $0\leqslant k\leqslant n$, and $\omega(l,m) = 1$ otherwise.*

It is easy to check that all the translations $T_{(l,m)}$, $(l,m)\in\mathbb{Z}^2$, are bounded. By Corollary 13, the action of $\mathbb{Z}^2$ on $L^p(\mathbb{Z}^2)$ for every $1\leqslant p<\infty$ is hypercyclic. But for fixed l,m, there are only finitely many points on the line $(l,m)\mathbb{Z}$ in which ω is different from 1; thus, the action of $S = \{T^n_{(l,m)}\}_{n\in\mathbb{N}}$ is not hypercyclic, by Proposition 12.

References

[1] Chen, Ch.-Ch. Hypercyclic weighted translations generated by non-torsion elements. *Arch. Math. (Basel)* **101** (2013), no. 2, 135–141.

[2] Conejero, J. A.; Peris, A. Hypercyclic translation C_0-semigroups on complex sectors. *Discrete Contin. Dyn. Syst.* **25** (2009), no. 4, 1195–1208.

[3] Conejero, J. A.; Müller, V.; Peris, A. Hypercyclic behaviour of operators in a hypercyclic C_0-semigroup. *J. Funct. Anal.* **244** (2007), no. 1, 342–348.

[4] de Vries, J. The local weight of an effective locally compact transformation group and the dimension of $L^2(G)$. *Colloq. Math.* 39 (1978), no. 2, 319–323.

[5] Desch, W.; Schappacher, W.; Webb, G. F. Hypercyclic and chaotic semigroups of linear operators. *Ergodic Theory Dyn. Syst.* **17** no. 4 (1997), 793–819.

[6] Edwards R. E. The stability of weighted Lebesgue spaces. *Trans. Amer. Math. Soc.* **93** (1959), 369–394.

[7] Feichtinger H. G. Gewichtsfunktionen auf lokalkompakten Gruppen, *Sitzber. Österr. Akad. Wiss. Abt. II,* **188**, no. 8–10 (1979), 451–471.
[8] Grosse-Erdmann, K.-G. Universal families and hypercyclic operators. *Bull. Amer. Math. Soc. (N.S.)* **36** (1999), no. 3, 345–381.
[9] Hewitt, E.; Ross, K.A. *Abstract harmonic analysis I, II.* Springer–Verlag, 3rd printing, 1997.
[10] Salas, H., *Hypercyclic weighted shifts, Trans. Amer. Math. Soc.* **347** (1995), 993–1004.
[11] Tam K. W. On measures with separable orbit. *Proc. Amer. Math. Soc.* **23** (1969), 409–411.

Evgeny Abakumov: University Paris-Est, 5 boulevard Descartes, 77454 Marne-la-Vallée, France
E-mail address: evgueni.abakoumov@u-pem.fr

Yulia Kuznetsova: University of Bourgogne Franche-Comté, 16 route de Gray, 25030 Besançon, France
E-mail address: yulia.kuznetsova@univ-fcomte.fr